Anaphylaxis Emergency Action Plan  Child's Name: Grade: Div: Birthdate:  School Name: School Address:					
School Name:	School Address:				
THIS PERSON HAS A POTENTIALLY LIFE THREATENING ALLERGY (ANAPHYLAXIS)			ACT QUICKLY. DO NOT WAIT FOR SYMPTOMS TO GET WORSE OR NEW SYMPTOMS TO BEGIN		
РНОТО	Allergy trigger(s):    Food(s):		<ul> <li>Give Epinephrine at the first sign of an anaphylactic reaction.</li> <li>Call 9-1-1</li> </ul>		
	☐ Insect Stings:		• Call Emergency Contact  Epinephrine is the first line medication for the emergency management of anaphylaxis. Antihistamines (e.g. Benedryl ™) or asthma medication should not be used to treat anaphylaxis.		
	Medication:  □ EpiPen Jr. (0.15mg) □ EpiPen Sr. (0.3mg)  Location: Expiry:				
AN ANAPHYLACTIC REACTION MAY HAVE THE FOLLOWING SIGNS AND SYMPTOMS:		EMERGENCY CONTACT INFO:			
<u>Face</u> : Hives, itching, swelling (lips, face, tongue) flushed face or body		Name	Relationship	Cell Phone	Other Phone
<u>Airway:</u> Difficulty breathing, swallowing or speaking, coughing, wheezing, change of voice, sneezing					
<b><u>S</u>tomach:</b> Stomach cramps, nausea, vomiting, diarrhea					
$\underline{\underline{\mathbf{T}}}$ otal body: Hives, itching, swelling, weakness, dizziness, loss of consciousness, anxiety, feeling of doom					
The undersigned parent/guardian authorizes any adult to administer emergency medication following the instructions outlined above to the above named person in the event of an anaphylactic reaction, as described above. This protocol has been recommended by a physician/NP. The plan will be shared with appropriate facility/school personnel to assist in responding in an Emergency. It is the parent/guardian's responsibility to advise the school about any changes to this plan.					
Parent/Guardian	Date Doctor	/NP Signature	 Date		

